




Massachusetts Eye and Ear Infirmary

PEDIATRIC ORDER SHEET

Room # Patient Weight: kg

Patient Allergies & Type of Reaction (nursing to verify)

PAIN SCALE



No Pain Mild Pain Moderate Pain Severe Pain

FLACC Pain Scale: For children under age 3 and for those who cannot self-report pain

Score each of the five categories from 0-2 and total the scores from each category, which results in a total score of 0-10

Table with 4 columns: FLACC Pain Scale, 0, 1, 2. Rows include Face, Legs, Activity, Cry, and Consolability with corresponding symptoms.

WARNINGS: DO NOT USE ABBREVIATIONS

- U, u
IU
Q.D, QD, qd., qd
Q.O.D, QOD, q.o.d., qod
Trailing zero (X.0mg)
Lack of leading zero (.xmg)
Any medication

ALL OTHER ORDERS

Assign to: MD

Admitting Diagnosis:

Status Post Procedure:

Pertinent Medical History: (for maintenance meds and nursing care)

Healthy Other

Allergies & Reactions:

Condition:

Vital Signs per floor routine Other

Diet NPO Reg Clear Soft Advanced as tolerated

Head position 45 Other

Activity as tolerated Other

Reinforce dressing PRN Suture line care with Bacitracin

Routine ice to: Neck OD OS

O2 Saturation spot check if < 94%, continuous O2 Sat monitoring

Continuous O2 Saturation monitoring

O2 @ % via

Special Notify Resident

T > 101.5 O2 < 94% HR > 120 or <70

Additional Orders on a separate order sheet

Discharge patient home when meets criteria with instructions, prescriptions and return visit information

Responsible physician(s) for additional issues:

Resident MD

Fellow MD

Attending MD

Date: Time:

Ordering MD Signature:

Ordering MD Printed Name:

Ordering MD Beeper /Phone:

MEDICATIONS

Date Time

WARNING: DO NOT USE D5W in PEDIATRIC PATIENTS

0-10 kg 100ml/kg/24 hours\*\*\*

11-20 kg 1000ml + 50ml/kg (for each kg > 10) / 24 hours\*\*\*

>20 kg 1500ml + 20ml/kg (for each kg > 20) / 24 hours\*\*\*

\*\*\*All results are divided by 24 = hourly rate\*\*\*

IV Fluid D5 LR @ ml/hr or @ ml/hr

Discontinue IV when tolerating PO fluids

If NO loading dose: Acetaminophen mg

(10-15mg/kg/dose) PO Q 4 hours PRN mild pain

If loading dose given Acetaminophen mg

(10-15mg/kg/dose) PO Q 6 hours PRN mild pain

1st post-op dosage at least 6 hours after initial dose

WARNING: Limit Acetaminophen to no more than 3 conventional dosage (15mg/kg) in 1st 24 hours

Roxicet oral liquid (Oxycodone 5mg/Acetaminophen 325mg per 5 ml) ml (based on Oxycodone 0.05 - 0.15 mg/kg/dose) PO Q 4 hours PRN moderate to severe pain

Acetaminophen & Codeine Oral Solution (300mg/30mg/12.5ml) ml (based on 0.5-1mg Codeine/kg/dose) PO Q4 hours PRN moderate to severe pain

WARNING: <12 years old Maximum Acetaminophen (including Roxicet) is 5 doses in 24 hours or 90 mg/kg/day

Ondansetron mg < 40kg 0.1mg/kg/dose IV x1 > 40kg 4 mg/dose IV x 1 prn nausea/vomiting

If patient continues nausea or vomiting use promethazine

Promethazine mg (0.25-1mg/kg/dose) PO Q 6 hours PRN for nausea/vomiting (maximum 25 mg/dose)

Promethazine mg (0.25-1mg/kg/dose) IV Q 6 hours PRN for nausea/vomiting (maximum 25 mg/dose)

WARNING: Promethazine is CONTRAINDICATED in Children < 24 months old

Antibiotics:

DO NOT WRITE IN THIS SPACE